

Some Aspects of the Wiener Index for Sun Graphs

Mohamed Amine Boutiche

USTHB, Faculty of Mathematics, LaROMaD Laboratory

BP 32 El Alia, 16111 Bab Ezzouar, Algiers, Algeria

mboutiche@usthb.dz

Abstract

The Wiener index $W(G)$ is the sum of distances of all pairs of vertices of the graph G . The Wiener polarity index $W_p(G)$ of a graph G is the number of unordered pairs of vertices u and v of G such that the distance $d_G(u, v)$ between u and v is 3. In this paper the Wiener and the Wiener polarity indices of sun graphs are computed. A relationship between those indices with some other topological indices are presented. Finally, we find the Hosoya (Wiener) polynomial for sun graphs.

Keywords. Wiener Index, Wiener polarity index, Sun Graphs.

AMS subject classifications. 05C85, 68R10, 68W05.

1 Introduction

In theoretical chemistry, distance based molecular structure descriptors are used for modeling physical, pharmacologic, biologic and others properties of chemical compounds [13, 14].

Let $G = (V, E)$ be a finite undirected connected graph. The distance between two vertices u and v in G , denoted by $d_G(u, v)$, is the length of a shortest path between u and v in G . A tree is a connected acyclic graph. Let $d_G(v)$ of a vertex v be the sum of all distances between v and all other vertices of G .

The Wiener index $W(G)$ of G is defined by

$$W(G) = \sum_{u, v \subseteq V(G)} d_G(u, v).$$

One can define the Wiener index also in a slightly different way

$$W(G) = \frac{1}{2} \sum_{v \in V(G)} d_G(v).$$

The Wiener polarity index $W_P(G)$ of a graph G of order n is defined as the number of unordered pairs of vertices u and v of G such that the distance $d_G(u, v)$ between u and v is 3. $W_P(G)$ was recently introduced and receive more attention. The authors in [12] demonstrated quantitative structure-property relationships in a series of acyclic and cycle-containing hydrocarbons by using $W_P(G)$. Hosoya [9] found a physical-chemical interpretation of $W_P(G)$. Recently, [5] gives all trees (resp. unicyclic graphs) of order n minimizing and maximizing the Wiener polarity. Some extremal properties on particular trees were given in [4, 2, 3, 11].

A k -sun graph ($k \geq 3$), is the graph on $2k$ vertices obtained from a clique $c_1, \dots, c_k$ on k vertices and an independent set $s_1, \dots, s_k$ on k vertices. The set of edges is defined by couples $s_i c_i, s_i c_{i+1}$, $1 \leq i < k$, and $s_k c_k, s_k c_1$, see figure 1.




Figure 1: A k -Sun graph

The paper is organized as follow, the Wiener and the Wiener polarity indices of sun graphs are computed in section 2. In section 3, a relationship between those indices with some other topological indices are presented. Finally, in section 4 we find the Hosoya (Wiener) polynomial for sun graphs.

2 Exact Wiener and Wiener Polarity Index of k -sun graphs

Let $G = (V, E)$ be a k -sun graph for $k \geq 3$, obtained from a clique $c_1, \dots, c_k$ on k vertices and an independent set $s_1, \dots, s_k$.

Theorem 1 *Let G be a k -sun graph ($k \geq 3$) then, $W(G) = k(4k - 5)$.*

Proof. We have by definition

$$\begin{aligned} W(G) &= \frac{1}{2} \sum_{v \in V(G)} d_G(v), \\ &= \frac{1}{2} \sum_{v \in U(G)} d_G(v) + \frac{1}{2} \sum_{v \in C(G)} d_G(v). \end{aligned}$$

Where $C(G)$ a clique $c_1, \dots, c_k$ on k vertices and $U(G)$ an independent set $s_1, \dots, s_k$ of G .

Let $U = \sum_{v \in U(G)} d_G(v)$ and $C = \sum_{v \in C(G)} d_G(v)$. Then, for a vertex u_i chosen arbitrarily, then

$$U = k \cdot (d(u_i, c_i) + d(u_i, c_{i+1}) + d(u_i, u_{i+1}) + d(u_i, u_{i-1}) + (k-2)d(u_i, c_j) + (k-3)d(u_i, u_j))$$

with $d(u_i, c_i) = 1$, $d(u_i, c_{i+1}) = 1$, $d(u_i, u_{i+1}) = 2$, $d(u_i, u_{i-1}) = 2$, $d(u_i, c_j) = 2$ and $d(u_i, u_j) = 3$. Hence, $U = k \cdot (5k - 7)$.

Moreover, for a vertex c_i chosen arbitrarily, we have $C = k \cdot ((k-1)d(c_i, c_j) + d(c_i, u_i) + d(c_i, u_{i+1}) + (k-2)d(c_i, u_j))$ with $d(c_i, c_j) = 1$, $d(c_i, u_i) = 1$, $d(c_i, u_{i+1}) = 1$, $d(c_i, u_j) = 2$. Then, $C = k \cdot (3k - 3)$, proving a result.

Theorem 2 Let G be a k -sun graph then $W_P(G) = \frac{k(k-3)}{2}$.

Proof. If $k = 3$, then $W_P(G) = 0$ since $\text{diam}(G) = 2$.

Let $G = (V, E)$ be a k -sun graph for $k \geq 4$.

Let compute $W_P(G) = |\{ \{u, v\} \mid d_G(u, v) = 3, u, v \in V \}|$.

All distances of all pairs of vertices $\{u, v\}$ such that $d_G(u, v) = 3$ are resumed below

If $u = c_i$ and $v = c_j$ then $d_G(u, v) \leq 1$ for $1 \leq i \leq k$, $1 \leq j \leq k$,

If $u = s_i$ and $v = c_i$ then $d_G(u, v) = 1$ for $1 \leq i \leq k$,

If $u = s_i$ and $v = c_{i+1}$ then $d_G(u, v) = 1$ for $1 \leq i \leq k$,

$d_G(u, v) = 2$, for all $u = s_i, v = c_j$, $j \neq i+1$, $i \neq k$ and $j \neq 1$, with $1 \leq i \leq k$, $1 \leq j \leq k$,

Finally, $d_G(u, v) = 3$ if and only if $u = s_i, v = s_j$, with $i \neq j$, $j \neq i+1$, $1 \leq i \leq k$, $1 \leq j \leq k$. $|\{ \{u, v\} \mid d_G(u, v) = 3 \}| = k - 3$, proving the result.

3 Relation between Wiener and Wiener Polarity index

The first Zagreb index $M_1(G)$, is defined as the summation of squares of the degrees of the vertices, and the second Zagreb index $M_2(G)$, is the sum of the products of the degrees of pairs of adjacent vertices of the graph G . These topological indices were introduced by Gutman and Trinajstić [6].

In [10] and for $d \geq 1$, authors define a generalization of Wiener polarity index as the number of unordered pairs of vertices $\{u, v\}$ of G such that the shortest distance between u and v is d .

$$W_d(G) = |\{\{u, v\} \mid d_G(u, v) = d, u, v \in V\}|.$$

So we have

$$W(G) = \sum_{d=1}^{\text{diam}(G)} W_d(G).$$

Lemma 3 [1] *Let G be a graph then*

$$W_p(G) \leq \frac{n(n-1)}{2} - \frac{1}{2}M_1(G),$$

with equality if $\text{diam}(G) = 3$.

Corollary 4 [1] *If $\text{diam}(G) \leq 3$ then $W(G) = \frac{3n(n-1)}{2} - \frac{1}{2}M_1(G) - m$.*

This corollary is valid for sun graphs since their diameter is ≤ 3 with $n = 2k$.

Corollary 5 *Let G be a sun graph then $W(G) = \frac{3n(n-1)}{2} - \frac{1}{2}M_1(G) - m$.*

Proposition 6 *Let G be a sun graph then $W(G) = n(n-1) + W_p(G) - m$.*

Proof. From corollary 1, 2 and lemma 1, we have

$$M_1(G) = n(n-1) + W_p(G) - m,$$

$$W(G) = \frac{3n(n-1)}{2} - \frac{1}{2}n(n-1) + W_p(G) - m - m,$$

$$\text{hence } W(G) = n(n-1) + W_p(G) - m.$$

4 The Hosoya (Wiener) polynomial of sun graphs

The Hosoya (Wiener) polynomial of graphs is defined by:

$$H(G, t) = \sum_{k=1}^D d(G, k)t^k.$$

The wiener index of a graph G can be determined as the first derivative of the polynomial $H(G, t)$ at $t = 1$, such that $W(G) = \sum_{k=1}^D k.t^k$

Theorem 7 Let G be a sun graph, then we have

$$H(G, t) = \frac{1}{2}k(k+3).t + k(k-1).t^2 + \frac{k(k-3)}{2}.t^3.$$

Proof.

We have $H(G, t) = \sum_{k=1}^3 d(G, k)t^k$, since G is a sun graph.

We have $d(G, 1) = |\{(u, v) \mid d(u, v) = 1\}|$, the pairs are $\{u_i, c_i\}_{i=1}^k$; $\{c_i, u_{i+1}\}_{i=1}^k$, and $\{c_i, c_j\}_{i,j=1}^k$. Then $d(G, 1) = k + k + \frac{k(k-1)}{2}$.

$d(G, 2) = |\{(u, v) \mid d(u, v) = 2\}|$, the pairs are $\{u_i, u_{i+1}\}_{i=1}^k$; $\{u_i, c_j\}_{i=1, j \neq i+1, i-1}^k$. Then $d(G, 2) = k + k(k-2) = k(k-1)$.

$d(G, 3) = |\{(u, v) \mid d(u, v) = 3\}|$, the pairs are $\{u_i, u_j\}_{i=1, j \neq i+1, i-1}^k$. Then $d(G, 3) = \frac{k(k-3)}{2}$, proving the result.

Remark 8 The wiener index of a sun graph G can be calculated in another way such as the first derivative of the polynomial $H(G, t)$ at $t = 1$, then

$$W(G) = \frac{1}{2}k(k+3) + 2k(k-1) + 3\frac{k(k-3)}{2} = k(4k-5)$$

Corollary 9 Let $G = S_3$ be the 3-sun graph, then $H(S_3, t) = 9.t + 6.t^2$.

References

- [1] A. Behmaram and H. Yousefi-Azari, *Further Results on Wiener Polarity Index of Graphs*, Iranian J. of Math. Chem. **2(1)** (2011), 67–70.
- [2] H. Deng, *On the extremal Wiener polarity index of chemical trees*, MATCH Commun. Math. Comput. Chem. **66** (2011), 305–314.
- [3] H. Deng and H. Xiao, *The maximum Wiener polarity index of trees with k pendants*, Appl. Math. Lett. **23** (2010), 710–715.
- [4] H. Deng, H. Xiao and F. Tang, *On the extremal Wiener polarity index of trees with a given diameter*, MATCH Commun. Math. Comput. Chem. **63** (2010), 257–264.
- [5] W. Du, X. Li and Y. Shi, *Algorithms and extremal problem on Wiener polarity index*, MATCH Commun. Math. Comput. Chem. **62** (2009), 235–244.
- [6] I. Gutman and N. Trinajstić, *Graph theory and molecular orbitals, Total φ -electron energy of alternant hydrocarbons*, Chem. phys. Lett **17** (1972), 535–538.
- [7] I. Gutman and E. Estrada, *Topological indices based on the line graph of the molecular graph*, J. Chem. Inf. Comput. Sci. **36** (1996), 541–543.

- [8] I. Gutman, L. Popovic, B.K. Mishra, M. Kaunar, E. Estrada and N. Guevara, *Application of line graphs in physical chemistry. Predicting surface tension of alkanes*, J. Serb. Chem. Soc. **62** (1997), 1025–1029.
- [9] H. Hosoya, *Mathematical and chemical analysis of Wiener’s polarity number*, in: D.H. Rouvray, R.B. King (Eds.), *Topology in Chemistry-Discrete Mathematics of Molecules*, Horwood, Chichester, 2002, p. 57.
- [10] A. Ilić and M. Ilić, *Generalizations of Wiener Polarity Index and Terminal Wiener Index*, Graphs and Combinatorics. **29** (2013), 1403–1416.
- [11] B. Liu, H. Hou and Y. Huang, *On the Wiener polarity index of trees with maximum degree or given number of leaves*, Comput. Math. Appl. **60** (2010), 2053–2057.
- [12] I. Lukovits and W. Linert, *Polarity-numbers of cycle-containing structures*, J. Chem. Inf. Comput. Sci. **38** (1998), 715–719.
- [13] N. Trinajstić, *Chemical Graph Theory*, CRC Press, Boca Raton, FL, 1993.
- [14] H. Wiener, *Structural determination of paraffin boiling points*, J. Amer. Chem. Soc. **69** (1947), 17–20.